

**Galzin**<sup>®</sup>  
(zinc acetate) capsules

*Wilson disease is a lifelong journey*



Experience a  
maintenance  
therapy that  
supports you  
every step of  
the way

Please see Indication and Important Safety Information below.



**Backed  
by science**



**Trusted  
by doctors**



**Personalized  
support for you**

#### **INDICATION**

GALZIN<sup>®</sup> (zinc acetate) is a prescription medicine used as a maintenance treatment of Wilson's disease in patients that were initially treated with a medication used to remove metallic deposits in the body.

#### **IMPORTANT SAFETY INFORMATION**

Always take GALZIN exactly as your doctor has directed.

**Please see Important Safety Information throughout and accompanying full Prescribing Information.**

## Your copper levels are under control. Now what?

You know that Wilson disease affects your body's ability to process and eliminate copper. When too much copper builds up in vital organs like the liver and brain, your symptoms can become more severe and can even lead to death. **That's why Wilson disease requires lifelong maintenance treatment.**

Once you've worked with your doctor to reduce the amount of copper in your body, the American Association for the Study of Liver Diseases (AASLD) **recommends zinc as a first-line maintenance therapy for Wilson disease.**

*Recommended*  
**FIRST-LINE  
MAINTENANCE  
THERAPY**

## Galzin® (zinc acetate) capsules can help keep the copper levels in your body in check.

*In a clinical study of 60 people living with Wilson disease*

 **91%** of copper balance tests (N=70) showed adequate control with the recommended dose of 50 mg of Galzin 3 times a day\*

\*Some people were prescribed different doses of Galzin at different frequencies.



**Galzin is the only FDA-approved zinc therapy for Wilson disease and offers what OTC supplements may not**



FDA-approved  
for 25+ years



Rigorous  
drug cGMP  
manufacturing  
standards



Demonstrated  
safety profile  
and results



Eton Cares®  
Patient Assistance  
Program

### IMPORTANT SAFETY INFORMATION (cont'd)

Do not take GALZIN if you are allergic to zinc acetate or any of its other ingredients.

You should take GALZIN on an empty stomach, at least one hour before or two to three hours after meals.

Capsules should be swallowed whole, not opened or chewed.

**Please see Important Safety Information throughout and accompanying full Prescribing Information.**

# Things you should know

**Galzin**<sup>®</sup>  
(zinc acetate) capsules

## Galzin has been used for decades

Gastric irritation (also known as gastritis) is the most commonly reported side effect of Galzin. About 10% of people experience a burning or disagreeable feeling in the stomach immediately after taking Galzin. For most people, it usually occurs with the morning dose and goes away within the first few days of therapy.

**Talk to your doctor about any discomfort you experience so they can help determine the best way for you to continue taking Galzin.**

**Please see additional Important Safety Information below.**

### Taking Galzin as prescribed is important for staying in control of your disease for the long term



Galzin is usually prescribed as **50 mg 3 times a day for adults.**



For patients who are compliant with therapy, **25 mg taken 3 times a day may be advisable.**

Galzin may also be prescribed as **25 mg 3 times a day for children 10 years of age or older and women who are pregnant.**



You should **take Galzin on an empty stomach**, at least 1 hour before or 2 to 3 hours after eating.



## IMPORTANT SAFETY INFORMATION (cont'd)

**Copper Deficiency:** Zinc acetate taken over long periods of time (i.e., months to years) may result in a loss of your body's ability to absorb copper and therefore very low levels of copper. Your doctor will monitor you for signs and/or symptoms of low levels of copper during treatment with GALZIN. Your treatment may be interrupted so your doctor can test for zinc levels.

**Gastric Ulcer:** Gastric (stomach) ulcers may develop with long-term use of zinc acetate. If you develop signs and/or symptoms of stomach ulcers during treatment with zinc acetate. Your doctor will have you discontinue GALZIN treatment.

**Please see Important Safety Information throughout and accompanying full Prescribing Information.**

# Eton Cares Program® - Comprehensive, personalized support every step of the way

eTon  
CARES

A prescription for Galzin® (zinc acetate) capsules automatically enrolls you in the Eton Cares Program®.

 **\$0 copay**  
for commercially eligible patients†

†Restrictions, limitations, and/or eligibility requirements may apply.

For questions or support, please contact us at our specialty pharmacy Anovo®:

 833-442-5946

 866-318-2990



Learn more at  
[www.galzin.com](http://www.galzin.com)

Your doctor sent your prescription to our specialty pharmacy Anovo.

## What's next?

- 1 Welcome Call**  
from your Care Coordinator to verify prescription, offer support, and answer questions.
- 2 Insurance Verification**  
to review and check coverage and benefits.
- 3 Financial Assistance Review**  
to explore copay programs, financing, or other options.
- 4 Medication Delivery**  
with instructions and specific product details.
- 5 Ongoing Support**  
with pharmacists for refills, check-ins, alerts, and 24/7 help

## IMPORTANT SAFETY INFORMATION (cont'd)

The most common adverse reactions are inflammation of the stomach lining, and increased levels of serum alkaline phosphatase (an enzyme that breaks down proteins), amylase (an enzyme that breaks starch into sugars), and lipase (an enzyme that helps the body digest fat) lasting from weeks to months suggesting pancreatitis (inflammation of the pancreas).

You are encouraged to report negative side effects of prescription drugs by contacting Eton Pharmaceuticals, Inc. at 1-855-224-0233 or the U.S. Food and Drug Administration (FDA) at [www.fda.gov/safety/medwatch](http://www.fda.gov/safety/medwatch) or call 1-800-FDA-1088.

Please see accompanying full Prescribing Information.